

MORC Family ATPases Required for Heterochromatin Condensation and Gene Silencing

Guillaume Moissiard,¹ Shawn J. Cokus,¹ Joshua Cary,¹ Suhua Feng,¹ Allison C. Billi,² Hume Stroud,¹ Dylan Husmann,¹ Ye Zhan,³ Bryan R. Lajoie,³ Rachel Patton McCord,³ Christopher J. Hale,¹ Wei Feng,⁴ Scott D. Michaels,⁴ Alison R. Frand,⁵ Matteo Pellegrini,^{1,6} Job Dekker,³ John K. Kim,² Steve Jacobsen^{1,5,6,7*}

¹Department of Molecular, Cell, and Developmental Biology, University of California at Los Angeles, Terasaki Life Sciences Building, 610 Charles Young Drive East, Los Angeles, CA 90095-723905, USA.

²Life Sciences Institute and Department of Human Genetics, University of Michigan, Ann Arbor, MI 48109, USA. ³Lazare Research Building 570N, Program in Systems Biology and Gene Function and Expression, University of Massachusetts Medical School, 364 Plantation Street, Worcester, MA 01605, USA.

⁴Department of Biology, Indiana University, Bloomington, IN 47405, USA. ⁵Department of Biological Chemistry, David Geffen School of Medicine, University of California Los Angeles, Los Angeles, CA 90095, USA. ⁶Eli and Edythe Broad Center of Regenerative Medicine and Stem Cell Research, University of California Los Angeles, Los Angeles, CA 90095, USA. ⁷Howard Hughes Medical Institute, University of California Los Angeles, Los Angeles, CA 90095, USA.

*To whom correspondence should be addressed. E-mail: jacobsen@ucla.edu

Transposable elements (TEs) and DNA repeats are commonly targeted by DNA and histone methylation to achieve epigenetic gene silencing. We isolated mutations in two *Arabidopsis* genes, *AtMORC1* and *AtMORC6*, which cause de-repression of DNA-methylated genes and TEs, but no losses of DNA or histone methylation. *AtMORC1* and *AtMORC6* are members of the conserved *Microrchidia* (MORC) adenosine triphosphatase (ATPase) family, predicted to catalyze alterations in chromosome superstructure. The *atmorc1* and *atmorc6* mutants show decondensation of pericentromeric heterochromatin, increased interaction of pericentromeric regions with the rest of the genome, and transcriptional defects that are largely restricted to loci residing in pericentromeric regions. Knockdown of the single MORC homolog in *Caenorhabditis elegans* also impairs transgene silencing. We propose that the MORC ATPases are conserved regulators of gene silencing in eukaryotes.

Gene silencing in the *Arabidopsis* genome is highly correlated with DNA methylation, which is found in three different cytosine contexts. Methylation of symmetric CG and CHG sites (in which H = A, T, or C) are mediated by DNA METHYLTRANSFERASE1 (MET1) and CHROMOMETHYLASE3 (CMT3), respectively, while CHH methylation is mainly catalyzed by DOMAINS REARRANGED METHYLTRANSFERASE2 (DRM2) (1). Silent loci are also enriched in the repressive histone H3 lysine 9 dimethylation mark (H3K9me2) (2, 3).

Suppressor of drm2 and *cmt3* (*SDC*) is a gene whose repression in most tissues depends on the redundant activities of DRM2 and CMT3 (4, 5). Hence, a loss of *SDC* silencing is observed in the *drm2 cmt3* double mutant but not in the *drm2* or *cmt3* single mutants. The *SDC* promoter carries seven tandem repeats, which recruit the DNA methylation machinery and cause transcriptional gene silencing. We engineered a GFP-based sensor construct controlled by the *SDC* promoter (fig. S1A). The *SDC::GFP* transgene behaves similarly to endogenous *SDC*, and GFP fluorescence is not detectable in wild-type, *drm2* or *cmt3* plants but is highly expressed in *drm2 cmt3* double mutant (Fig. 1A).

We carried out ethyl methanesulfonate (EMS) mutagenesis screens in wild type or *cmt3* backgrounds for mutants showing *SDC::GFP* overexpression, and identified the wt #67, *cmt3* #7 and *cmt3* #49 mutants

(Fig. 1, A and B). Mapping experiments using bulk segregant analysis coupled to deep genome re-sequencing indicated that *cmt3* #7 contained a mutation in *At4g36290* (*AtMORC1*), previously also named *COMPROMISED RECOGNITION OF TCV-1* (*CRT1*) (6, 7), while wt #67 and *cmt3* #49 both contained mutations in *At1g19100* (*AtMORC6*) (7) (figs. S1B, S2, and S3A). An *atmorc1* allele was previously reported to show reduced resistance to the TCV virus (6, 7), suggesting that *AtMORC1* is also involved in viral resistance, in addition to its role in gene silencing described in this study; whereas mutations in *AtMORC6* have not been described. To ensure that *atmorc1* and *atmorc6* mutations were those responsible for the loss of *SDC* silencing, we isolated knock-out T-DNA insertion lines *atmorc1-4* and *atmorc6-3*, and confirmed *SDC* overexpression in these two mutant alleles (fig. S3B, C and D). Genetic complementation crosses between the recessive EMS and T-DNA mutants confirmed *AtMORC1* and *AtMORC6* as the mutated genes responsible for *SDC::GFP* activation in the three EMS lines (fig. S3E). Therefore, #7, #67 and #49 were renamed *atmorc1-3*, *atmorc6-1* and *atmorc6-2*, respectively.

Using RNA sequencing (RNA-seq) (8), we found that the majority of RNAs significantly affected in the *atmorc1* and *atmorc6* mutants showed up-regulation, and many of these TEs belong to various transposon super families including, among others, the

LTR/Gypsy, LTR/Copia, DNA/MuDR and DNA/Harbinger families (Fig. 1C and D, fig. S4A, table S1). The expression defects in the *atmorc1* and *atmorc6* mutants were very similar, with all but two of the transposons upregulated in *atmorc1* also upregulated in *atmorc6* (fig. S4B). Protein-coding genes overexpressed in the *atmorc1* and *atmorc6* EMS and T-DNA mutants included endogenous *SDC* (table S2). There was a high degree of overlap between the genes upregulated in *atmorc1* and *atmorc6* (fig. S4C), most of them corresponding to DNA-methylated and silenced loci, derepressed in both *atmorc1* and *atmorc6* (fig. S4D and E). We also performed RNA-seq in the *atmorc1 atmorc6* double mutant and found a very similar set of genes and transposons upregulated, with only a few genes upregulated in the double mutant that were not upregulated in each of the single mutants (table S3), suggesting that *AtMORC1* and *AtMORC6* may act together to enforce gene silencing.

Whole genome bisulfite sequencing (BS-seq) (9) revealed that DNA methylation levels in all sequence contexts were unaltered in *atmorc1* or *atmorc6* relative to wild type at TEs upregulated in *atmorc1* or *atmorc6* (Fig. 2A and B) nor were there any bulk alterations in protein-coding genes or TEs in the genome (fig. S5A and B). In addition, analyses at the pericentromeric satellite CEN180 repeats and five loci upregulated in *atmorc1* and *atmorc6* showed that the DNA methylation patterns in *atmorc1-4* and *atmorc6-3* were similar to wild type (Fig. 2C and D). ChIP-





A**B****C****D**

Fig. 1. Mutations of two MORC homologs induce *SDC::GFP* and TE overexpression. **(A)** Wild type (wt), *drm2* mutant and *cmt3* mutant plants carrying *SDC::GFP* showing no GFP fluorescence under UV light (insets show each plant under white light) and *drm2 cmt3* double mutant and EMS-mutagenized lines wt #67, *cmt3* #49, and *cmt3* #7 plants showing strong GFP fluorescence. **(B)** Western blot using anti-GFP antibody confirms *SDC::GFP* overexpression in the EMS mutants. Coomassie staining of the large Rubisco subunit (*rbcL*) is used as loading control. **(C)** Number of TEs overexpressed in *atmorc1* and *atmorc6* and classified by superfamily. For each mutant, only TEs with a ≥ 4 fold increase in both the EMS and T-DNA alleles over wild type and with a p -value ≤ 0.05 are represented. **(D)** Relative fold increase of four TE transcripts in *atmorc1-4* and *atmorc6-3* over wild type assayed by Real-Time quantitative PCR (RT-qPCR) and normalized to *ACTIN7*. Errors bars indicate standard deviation based on three independent biological replicates.

seq analyses of H3K9me2 also did not reveal any changes in the *atmorc1* or *atmorc6* mutants at *SDC* or other upregulated locations (fig. S6A and B). Finally, small RNA sequencing analyses showed that elements up-regulated in *atmorc1* and *atmorc6* mutants were enriched in siRNAs, but these siRNA levels did not change in the mutants (fig. S7). Thus *AtMORC1* and *AtMORC6* are not required to maintain DNA methylation, H3K9me2, or siRNAs, suggesting that *AtMORC1* and *AtMORC6* are likely to either act downstream of DNA methylation or enforce silencing by a novel mechanism.

AtMORC1 and *AtMORC6* are homologs of mouse *Microrchidial* (*MORC1*) (10, 11), and contain Gyrase, Hsp90, Histidine Kinase and MutL (GHKL) and S5 domains, together comprising an ATPase module (6) in addition to a putative C terminal coiled-coil domain (fig. S1B). The EMS mutations found in *atmorc1-3*, *atmorc6-1* and *atmorc6-2* alleles all introduced premature stop codons within the GHKL domain (fig. S1B).

Because of the similarity of *AtMORC1* and *AtMORC6* to ATPases involved in manipulating chromatin superstructure (12), these proteins may affect gene silencing through higher order compaction of methylated and silent chromatin. In wild type nuclei, pericentromeric heterochromatin forms densely staining nuclear bodies called chromocenters that localize to the nuclear periphery (13). We observed decondensation of chromocenters in the *atmorc1* and *atmorc6* mutants (as well as in *atmorc1 atmorc6* double mutant) (fig. S8-11), and also found that loci transcriptionally de-repressed in the mutants mostly localized to pericentromeric heterochromatin (fig. S12, tables S1 and S3). To directly examine whole genome chromatin interactions, we performed Hi-C analyses in wild type and *atmorc6-1* (14). Consistent with previous cytological studies (13), the wild type genome showed interactions between telomeres as well as between euchromatic regions on the same chromosome arm (Fig. 3A). In contrast, pericentromeric heterochromatin regions interacted very weakly with the rest of the genome, consistent with their compaction in chromocenters (Fig. 3A). While *atmorc6-1* showed a roughly similar chromatin architecture (fig. S13), plotting the differences between mutant and wild type showed that *atmorc6-1* shows an increase in interactions between the pericentromeric regions of all chromosomes with the euchromatic arms of all chromosomes, and a corresponding depletion of interactions of euchromatic arms with themselves. Since the analysis reports relative changes, with the sum of differences set to zero, the most likely interpretation of these findings is that pericentromeric regions interact more strongly with the euchromatic arms in *atmorc6-1*, although we cannot exclude that the mutant also has effects on the euchromatic arms (Fig. 3B). This interpretation is consistent with the cytological observations showing that chromocenters expand out into a larger area of the nucleus in the mutants (fig. S8). We also found, using complementing myc-tagged transgenes, that *AtMORC1* and *AtMORC6* proteins formed small nuclear bodies that were usually adjacent to, but not within chromocenters (Fig. 3C and figs. S14-15). These results are all consistent with a model in which *AtMORC1* and *AtMORC6* enforce compaction and gene silencing of pericentromeric heterochromatin, although it is also possible that changes in chromatin and gene expression in the mutant secondarily lead to the observed changes in chromatin compaction. Mutation of the plant specific *MOM1* gene has also been shown to affect gene silencing but not DNA methylation in *Arabidopsis*, however *mom1* mutants do not show chromocenter decondensation and therefore are likely to act via a different mechanism (15, 16).

A single MORC homolog, *morc-1*, is present in the worm *Caenorhabditis elegans*, which is devoid of DNA methylation (17). To test if the *C. elegans morc-1* (ZC155.3) is involved in gene silencing, we performed RNAi-mediated knock-down of *morc-1* in the *eri-1* sensitized background, in which a GFP transgene is silenced in most of the worm seam cells (Fig. 3D) (18). *morc-1* depleted worms showed GFP reactivation similar to worms depleted of *rde-4*, an essential component of gene




Fig. 2. DNA methylation is not impaired in *atmorc1* and *atmorc6* mutants. (A and B) Metaplot analyses showing DNA methylation level in *atmorc1-4*, *atmorc6-3* and wild type for the set of TEs upregulated in *atmorc1-4* (A) and *atmorc6-3* (B). The gray vertical lines mark the boundaries between 1 kilobase upstream and downstream regions of TEs. (C) Southern blot analyses assaying CG methylation level at CEN180 repeats using *Hpa*II-treated genomic DNAs. m, methylated; u, unmethylated. *met1-3* genomic DNA is used as positive control for loss of CG methylation (23). (D) Percent DNA methylation at *SDC* and 4 TEs overexpressed in *atmorc1-4* and *atmorc6-3* assayed by bisulfite sequencing. 24 clones were analyzed for each individual analysis.

silencing in *C. elegans* (Fig. 3D) (19). These results suggest that MORCs may play an ancient and conserved role in gene silencing. In addition, the observation that *morc-1* is required for gene silencing in *C. elegans* reinforces our view that MORCs in *Arabidopsis* are enforcing silencing by a mechanism that may not be directly linked with DNA methylation. It is interesting to note that the phenotype of the *Morc1*-knockout mouse resembles *Miw12*- and *Dnmt3L*-knockout mouse phenotypes, showing male-specific meiotic defects during spermatogenesis (10, 20-22). *Miw12* and *Dnmt3L* are both required for TE silencing, and it is possible that *Morc1* might be involved in transposon silencing in mammals as well. We propose that MORC family ATPases acts to regulate chromatin architecture and gene silencing in a wide variety of eukaryotes.




Fig. 3. AtMORC1 and AtMORC6 are required for maintenance of chromatin architecture, form nuclear bodies near chromocenters, and *morc-1* is involved in gene silencing in *C. elegans*. **(A)** Interaction matrix of the wild type *Arabidopsis* genome from Hi-C analysis. Positions along the 5 chromosomes are shown from left to right and top to bottom, and each pixel represents interactions from uniquely mapping paired end reads in 200 kilobase bins. Black bars and circles mark the positions of the pericentromeric and telomeric regions, respectively. Light grey regions represent areas masked out due to problematic mapping. Black bars show separation between chromosomes. **(B)** Difference plot showing enrichment of Hi-C interactions in *atmorc6-1* in red and interactions depleted in *atmorc6-1* in blue. **(C)** Anti-Myc immunostaining showing localization of pAtMORC6::AtMORC6-Myc and pAtMORC1::AtMORC1-Myc in nuclear bodies adjacent to chromocenters. AtMORC1 and AtMORC6 showed an average of 2.0 +/- 1.0 (average plus or minus standard deviation) and 2.5 +/- 1.2 bodies per chromocenter, respectively. DAPI staining shows chromocenter location. Bottom panels are merges. **(D)** A silenced seam cell-specific GFP transgene in the *eri-1* (*mg366*) sensitized background is overexpressed in worms fed with bacteria expressing double-stranded RNA targeting *morc-1* or *rde-4* but not in worms fed with bacteria expressing a control empty vector. Results are representative of five independent replicates.

References and Notes

1. J. A. Law, S. E. Jacobsen, Establishing, maintaining and modifying DNA

methylation patterns in plants and animals. *Nat. Rev. Genet.* **11**, 204 (2010). [doi:10.1038/nrg2719](https://doi.org/10.1038/nrg2719) [Medline](#)

2. J. P. Jackson, A. M. Lindroth, X. Cao, S. E. Jacobsen, Control of CpNpG DNA methylation by the KRYPTONITE histone H3 methyltransferase. *Nature* **416**, 556 (2002). [doi:10.1038/nature731](https://doi.org/10.1038/nature731) [Medline](#)

3. F. Malagnac, L. Bartee, J. Bender, An Arabidopsis SET domain protein required for maintenance but not establishment of DNA methylation. *EMBO J.* **21**, 6842 (2002). [doi:10.1093/emboj/cdf687](https://doi.org/10.1093/emboj/cdf687) [Medline](#)

4. X. Zhang *et al.*, Genome-wide high-resolution mapping and functional analysis of DNA methylation in Arabidopsis. *Cell* **126**, 1189 (2006). [doi:10.1016/j.cell.2006.08.003](https://doi.org/10.1016/j.cell.2006.08.003) [Medline](#)

5. I. R. Henderson, S. E. Jacobsen, Tandem repeats upstream of the Arabidopsis endogene SDC recruit non-CG DNA methylation and initiate siRNA spreading. *Genes Dev.* **22**, 1597 (2008). [doi:10.1101/gad.1667808](https://doi.org/10.1101/gad.1667808) [Medline](#)

6. H. G. Kang, J. C. Kuhl, P. Kachroo, D. F. Klessig, CRT1, an Arabidopsis ATPase that interacts with diverse resistance proteins and modulates disease resistance to turnip crinkle virus. *Cell Host Microbe* **3**, 48 (2008). [doi:10.1016/j.chom.2007.11.006](https://doi.org/10.1016/j.chom.2007.11.006) [Medline](#)

7. H. G. Kang *et al.*, Endosome-associated CRT1 functions early in resistance gene-mediated defense signaling in Arabidopsis and tobacco. *Plant Cell* **22**, 918 (2010). [doi:10.1105/tpc.109.071662](https://doi.org/10.1105/tpc.109.071662) [Medline](#)

8. Z. Wang, M. Gerstein, M. Snyder, RNA-Seq: a revolutionary tool for transcriptomics. *Nat. Rev. Genet.* **10**, 57 (2009). [doi:10.1038/nrg2484](https://doi.org/10.1038/nrg2484) [Medline](#)

9. S. J. Cokus *et al.*, Shotgun bisulphite sequencing of the Arabidopsis genome reveals DNA methylation patterning. *Nature* **452**, 215 (2008). [doi:10.1038/nature06745](https://doi.org/10.1038/nature06745) [Medline](#)

10. M. L. Watson *et al.*, Identification of *morc* (microorchidia), a mutation that results in arrest of spermatogenesis at an early meiotic stage in the mouse. *Proc. Natl. Acad. Sci. U.S.A.* **95**, 14361 (1998). [doi:10.1073/pnas.95.24.14361](https://doi.org/10.1073/pnas.95.24.14361) [Medline](#)

11. N. Inoue *et al.*, New gene family defined by MORC, a nuclear protein required for mouse spermatogenesis. *Hum. Mol. Genet.* **8**, 1201 (1999). [doi:10.1093/hmg/8.7.1201](https://doi.org/10.1093/hmg/8.7.1201) [Medline](#)

12. L. M. Iyer, S. Abhiman, L. Aravind, MutL homologs in restriction-modification systems and the origin of eukaryotic MORC ATPases. *Biol. Direct* **3**, 8 (2008). [doi:10.1186/1745-6150-3-8](https://doi.org/10.1186/1745-6150-3-8) [Medline](#)

13. P. Fransz, J. H. De Jong, M. Lysak, M. R. Castiglione, I. Schubert, Interphase chromosomes in Arabidopsis are organized as well defined chromocenters from which euchromatin loops emanate. *Proc. Natl. Acad. Sci. U.S.A.* **99**, 14584 (2002). [doi:10.1073/pnas.212325299](https://doi.org/10.1073/pnas.212325299) [Medline](#)

14. E. Lieberman-Aiden *et al.*, Comprehensive mapping of long-range interactions reveals folding principles of the human genome. *Science* **326**, 289 (2009). [doi:10.1126/science.1181369](https://doi.org/10.1126/science.1181369) [Medline](#)

15. P. Amedeo, Y. Habu, K. Afsar, O. Mittelsten Scheid, J. Paszkowski, Disruption of the plant gene MOM releases transcriptional silencing of methylated genes. *Nature* **405**, 203 (2000). [doi:10.1038/35012108](https://doi.org/10.1038/35012108) [Medline](#)

16. A. V. Probst, P. F. Fransz, J. Paszkowski, O. Mittelsten Scheid, Two means of transcriptional reactivation within heterochromatin. *Plant J.* **33**, 743 (2003). [doi:10.1046/j.1365-3113X.2003.01667.x](https://doi.org/10.1046/j.1365-3113X.2003.01667.x) [Medline](#)

17. V. J. Simpson, T. E. Johnson, R. F. Hammen, *Caenorhabditis elegans* DNA does not contain 5-methylcytosine at any time during development or aging.

- Nucleic Acids Res.* **14**, 6711 (1986). doi:10.1093/nar/14.16.6711 [Medline](#)
18. S. Kennedy, D. Wang, G. Ruvkun, A conserved siRNA-degrading RNase negatively regulates RNA interference in *C. elegans*. *Nature* **427**, 645 (2004). doi:10.1038/nature02302 [Medline](#)
 19. H. Tabara *et al.*, The rde-1 gene, RNA interference, and transposon silencing in *C. elegans*. *Cell* **99**, 123 (1999). doi:10.1016/S0092-8674(00)81644-X [Medline](#)
 20. M. A. Carmell *et al.*, MIWI2 is essential for spermatogenesis and repression of transposons in the mouse male germline. *Dev. Cell* **12**, 503 (2007). doi:10.1016/j.devcel.2007.03.001 [Medline](#)
 21. D. Bourc'his, T. H. Bestor, Meiotic catastrophe and retrotransposon reactivation in male germ cells lacking Dnmt3L. *Nature* **431**, 96 (2004). doi:10.1038/nature02886 [Medline](#)
 22. A. A. Aravin *et al.*, A piRNA pathway primed by individual transposons is linked to de novo DNA methylation in mice. *Mol. Cell* **31**, 785 (2008). doi:10.1016/j.molcel.2008.09.003 [Medline](#)
 23. M. W. Kankel *et al.*, Arabidopsis MET1 cytosine methyltransferase mutants. *Genetics* **163**, 1109 (2003). [Medline](#)
 24. N. Kleinboelting, G. Huep, A. Kloetgen, P. Viehoveer, B. Weisshaar, GABI-Kat SimpleSearch: new features of the Arabidopsis thaliana T-DNA mutant database. *Nucleic Acids Res.* **40**, (Database issue), D1211 (2012). doi:10.1093/nar/gkr1047 [Medline](#)
 25. S. J. Clough, A. F. Bent, Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant J.* **16**, 735 (1998). doi:10.1046/j.1365-313x.1998.00343.x [Medline](#)
 26. J. A. Law, A. A. Vashisht, J. A. Wohlschlegel, S. E. Jacobsen, SHH1, a homeodomain protein required for DNA methylation, as well as RDR2, RDM4, and chromatin remodeling factors, associate with RNA polymerase IV. *PLoS Genet.* **7**, e1002195 (2011). doi:10.1371/journal.pgen.1002195 [Medline](#)
 27. M. V. Greenberg *et al.*, Identification of genes required for de novo DNA methylation in Arabidopsis. *Epigenetics* **6**, 344 (2011). doi:10.4161/epi.6.3.14242 [Medline](#)
 28. B. Langmead, C. Trapnell, M. Pop, S. L. Salzberg, Ultrafast and memory-efficient alignment of short DNA sequences to the human genome. *Genome Biol.* **10**, R25 (2009). doi:10.1186/gb-2009-10-3-r25 [Medline](#)
 29. A. Mortazavi, B. A. Williams, K. McCue, L. Schaeffer, B. Wold, Mapping and quantifying mammalian transcriptomes by RNA-Seq. *Nat. Methods* **5**, 621 (2008). doi:10.1038/nmeth.1226 [Medline](#)
 30. Y. Benjamini, Y. Hochberg, *J. R. Stat. Soc., B* **57**, 12 (1995).
 31. P. Y. Chen, S. J. Cokus, M. Pellegrini, BS Seeker: precise mapping for bisulfite sequencing. *BMC Bioinformatics* **11**, 203 (2010). doi:10.1186/1471-2105-11-203 [Medline](#)
 32. J. Lee *et al.*, Analysis of transcription factor HY5 genomic binding sites revealed its hierarchical role in light regulation of development. *Plant Cell* **19**, 731 (2007). doi:10.1105/tpc.106.047688 [Medline](#)
 33. C. Lu, B. C. Meyers, P. J. Green, Construction of small RNA cDNA libraries for deep sequencing. *Methods* **43**, 110 (2007). doi:10.1016/j.ymeth.2007.05.002 [Medline](#)
 34. Y. Zhang *et al.*, Spatial organization of the mouse genome and its role in recurrent chromosomal translocations. *Cell* **148**, 908 (2012). doi:10.1016/j.cell.2012.02.002 [Medline](#)
 35. W. J. Soppe *et al.*, DNA methylation controls histone H3 lysine 9 methylation and heterochromatin assembly in Arabidopsis. *EMBO J.* **21**, 6549 (2002). doi:10.1093/emboj/cdf657 [Medline](#)
 36. J. K. Kim *et al.*, Functional genomic analysis of RNA interference in *C. elegans*. *Science* **308**, 1164 (2005). doi:10.1126/science.1109267 [Medline](#)
 37. R. S. Kamath *et al.*, Systematic functional analysis of the *Caenorhabditis elegans* genome using RNAi. *Nature* **421**, 231 (2003). doi:10.1038/nature01278 [Medline](#)
 38. Y. Y. Polosina, C. G. Cupples, Wot the 'L-Does MutL do? *Mutat. Res.* **705**, 228 (2010). doi:10.1016/j.mrrev.2010.07.002 [Medline](#)
 39. Y. Mimura, K. Takahashi, K. Kawata, T. Akazawa, N. Inoue, Two-step colocalization of MORC3 with PML nuclear bodies. *J. Cell Sci.* **123**, 2014 (2010). doi:10.1242/jcs.063586 [Medline](#)
 40. T. Hirano, The ABCs of SMC proteins: two-armed ATPases for chromosome condensation, cohesion, and repair. *Genes Dev.* **16**, 399 (2002). doi:10.1101/gad.955102 [Medline](#)
 41. M. E. Blewitt *et al.*, SmcHDI, containing a structural-maintenance-of-chromosomes hinge domain, has a critical role in X inactivation. *Nat. Genet.* **40**, 663 (2008). doi:10.1038/ng.142 [Medline](#)
 42. I. Ausin, T. C. Mockler, J. Chory, S. E. Jacobsen, IDN1 and IDN2 are required for de novo DNA methylation in Arabidopsis thaliana. *Nat. Struct. Mol. Biol.* **16**, 1325 (2009). doi:10.1038/nsmb.1690 [Medline](#)
 43. T. Kanno *et al.*, A structural-maintenance-of-chromosomes hinge domain-containing protein is required for RNA-directed DNA methylation. *Nat. Genet.* **40**, 670 (2008). doi:10.1038/ng.119 [Medline](#)
 44. G. Böhmendorfer *et al.*, GMI1, a structural-maintenance-of-chromosomes-hinge domain-containing protein, is involved in somatic homologous recombination in Arabidopsis. *Plant J.* **67**, 420 (2011). doi:10.1111/j.1365-313X.2011.04604.x [Medline](#)
- Acknowledgments:** We thank M. Akhavan for sequencing, L. Goddard and L. Iruela-Arispe for assistance with confocal microscopy, and P. Frasz and I. Schubert for helpful discussions. S.F. is a Special Fellow of the Leukemia and Lymphoma Society. J.C. is supported by the Ruth L. Kirschstein National Research Service Award GM007185. R.P.M. is supported by the National Institute of General Medical Sciences (grant F32GM100617), and J.D. is supported by a W.M. Keck Foundation Distinguished Young Scholar in Medical Research Award. Research in the Jacobsen, Kim, Michaels, and Dekker laboratories was supported by NIH grants GM60398, GM088565, GM075060, and HG003143, respectively. Sequencing files have been deposited at GEO (accession code GSE37644). The authors declare no competing financial interests. S.E.J. is an investigator of the Howard Hughes Medical Institute. Correspondence and requests for materials should be addressed to S.E.J. (jacobsen@ucla.edu).
- Supplementary Materials**
www.sciencemag.org/cgi/content/full/science.1221472/DC1
 Materials and Methods
 Figs. S1 to S15
 Tables S1 to S4
 References (24–44)
- 05 March 2012; accepted 24 April 2012
 Published online 03 May 2012; 10.1126/science.1221472
 10.1126/science.1221472